

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO

Cloning, expression, and purification of recombinant protein MPT-64 from a virulent strain of *Mycobacterium bovis* in a prokaryotic system

Maryam Mohammadi Tashakkori^a, Majid Tebianian^{b,*}, Mohammad Tabatabaei^a, Nader Mosavari^a

^a Department of Pathobiology, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

^b Department of Biotechnology, Razi Vaccine and Serum Research Institute, Karaj, Iran

ARTICLE INFO

Article history:

Received 26 September 2016

Accepted 1 October 2016

Available online xxxx

Keywords:

Cloning

MPT-64 protein

Mycobacterium bovis

Recombinant protein

ABSTRACT

Objective: Tuberculosis (TB) is a zoonotic infectious disease common to humans and animals that is caused by the rod-shaped acid-fast bacterium *Mycobacterium bovis*. Rapid and sensitive detection of TB is promoted by specific antigens. Virulent strains of the TB complex from *M. bovis* contain 16 regions of difference (RD) in their genome that encode important proteins, including major protein of *Mycobacterium Tuberculosis* 64 (MBT-64, which is a primary immune-stimulating antigen encoded by RD-2. In this study, we cloned, expressed, and purified MPT-64 as a potent *M. bovis* antigen in a prokaryotic system for use in future diagnostic studies.

Methods: The antigenic region of the *Mpt64* gene was investigated by bioinformatics methods, cloned into the PQE-30 plasmid, and expressed in *Escherichia coli* M15 cells, followed by isopropyl β -D-1-thiogalactopyranoside induction. The expressed protein was analyzed sodium dodecyl sulfate polyacrylamide gel electrophoresis and purified using a nickel-affinity column. Biological activity was confirmed by western blot using specific antibodies.

Results: Our data verified the successful cloning of the *Mpt64* gene (687-bp segment) via the expression vector and purification of recombinant MPT-64 as a 24-kDa protein.

Conclusion: These results indicated successful expression and purification of recombinant MPT-64 protein in a prokaryotic system. This protein can be used for serological diagnosis, improved detection of pathogenicity and non-pathogenicity between infected cattle, and for verification of suspected cases of bovine TB.

Conflicts of interest

The authors have no conflicts of interest to declare.

* Corresponding author at: PhD of Immunology, Razi Vaccine & Serum Research Institute, P.O.BOX: 31975/148, Karaj, Iran. Tel.: +98 261 4570038 46 (2216); fax: +98 261 4552194.

E-mail address: tebianianm@hotmail.com (M. Tebianian).

Peer review under responsibility of Asian African Society for Mycobacteriology.

<http://dx.doi.org/10.1016/j.ijmyco.2016.10.027>